
Subject: Re: Doubly Substituted Reactants for Combinatorial Library
Posted by [nbehrnd](#) on Thu, 24 Jun 2021 04:46:08 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear sn,

from the documentation and some limited testing with DW I infer that this (currently) is not yet possible this way. Perhaps a misunderstanding by mine, but I infer «any atom number» is about «at this position, the atom may be a carbon, a nitrogen, an oxygen, etc», rather than the number of atoms (like let there be one atom, or two, three, etc.). On the other hand, one may prepare a list of this e.g., diamines separated by one, two, three methylene groups in advance. Past mapping the reaction in the tab «generic reaction», the subsequent tab «reactants» then allows to add them from a e.g., from a .dwar, .sdf, .mol2 file.

As an example, a bis-substitution like defined by


yields a list like

To obtain the special permutation «use the same acid on the left and on the right hand amine» would then justify to load and re-use such a pre-defined list to cover the chemical space efficiently.

Norwid

File Attachments

1) [setup.png](#), downloaded 893 times





2) [example.png](#), downloaded 930 times

File Edit Data Chemistry D

Generic Reaction Reactants

Table

	Product	Reactants
1		1
2		1


Structure

Suggest... <

Suggest... <

Suggest... <