

I will try to give at least a short explanation that summarizes the calculation:

Molecular flexibility values range from 0.0 (rigid) to 1.0 (completely flexible).
Four steps are needed for its calculation:

- All rotatable bonds are determined. These are the only considered contributors to flexibility. Bending or stretching the molecule or individual bonds is not considered. Double or triple bonds and bonds within aromatic rings or within 3-, 4- or 5-membered rings are not considered flexible. All remaining single bonds are considered rotatable, if both atoms are sp² or sp³ and carry at least one more non-hydrogen neighbor. For chains of conjugated triple bonds the following applies: If at least one terminal sp²/sp³ atom has no external neighbor, then no single bond is considered rotatable. Otherwise that terminal single bond connecting the smaller substituent is considered the only rotatable bond of the linear atom strand.
- Dihedral angle changes of a bond in the center of a molecule have a much larger effect on the overall molecule shape than the torsion change of an outer substituent bond. Therefore an influence factor is calculated for every rotatable bond. It also considers the limited degree of freedom of ring bond rotation:
 - for ring bonds: factor=0.33, since ring bonds cannot be changed without typically affecting two other rings
 - other bonds: factor= $\sqrt{2.0 * \text{smallerSideNonHydrogenAtomCount} / \text{moleculeNonHydrogenAtomCount}}$
- The flexibility contribution of every rotatable bond is determined from statistical torsion data of similar bonds in the Crystallography Open Database (COD). For instance C-O-C-C (C-O bond) is considered much more flexible than the central bond in a C-C-C-C chain, which has a strong preference towards the anti-conformation. For every rotatable bond a characterizing substructure is generated from its local environment. This substructure consists of the two bond atoms, their direct neighbour atoms and the next shell of neighbour atoms. It includes all connecting bond orders and various properties, e.g. whether atoms are in a ring, are aromatic, the stereo configuration, etc. Then canonical representation of the bond's characterizing substructure is built. The same mechanism was used before to characterize all rotatable bonds of all organic structures of the COD. Now using the canonical key DataWarrior gets a torsion angle distribution curve of equal bond environments

from all organic structures of the COD. The number of peaks, their heights and their widths are now used to generate the bond specific flexibility value. Frequency distributions with wide and multiple distribution maxima of similar heights receive local flexibility values close to 1.0 while histograms with one narrow single peak are close to 0.0.

- Finally, from the number of all molecule bonds, the number of rotatable bonds, their specific flexibility values, and their geometry weighting factor an overall flexibility value is calculated with a non-linear incremental approach.

For more details I suggest to check the source code in `com.actelion.research.chem.conf.MolecularFlexibilityCalculator`
