

Subject: Pentavalent carbonyl issue (urea formation?)

Posted by [padster](#) on Thu, 22 Jun 2023 19:14:13 GMT

[View Forum Message](#) <> [Reply to Message](#)


Hi, I am trying to create a combinatorial library (macrocyclic peptide) in DataWarrior. The issue I run into however is that every amide bond forms correctly with the next amino acid, but for some reason it also forms a bond (red bond in image) between the N and C=O of the same amino acid, creating a urea-like motif (refer to image attached). If the red bond is deleted, it forms the correct structure. Curiously, this issue does not happen when running DW on Mac. Is there a way to fix this on Windows so that it runs correctly as well, or is there some sort of setting that changes this? I reinstalled DW, but still run into the same issue.

Macrocycle:


Reaction:

File Attachments

1) [Screenshot_1.png](#), downloaded 1022 times



2) [Screenshot_2.png](#), downloaded 994 times



Subject: Re: Pentavalent carbonyl issue (urea formation?)

Posted by [thomas](#) on Fri, 23 Jun 2023 07:26:41 GMT

[View Forum Message](#) <> [Reply to Message](#)

Can you post the reaction as .rxn file, please? Or better a small macro containing the complete combinatorial library task with reaction and one real reactant for each generic reactant? This way I could directly reproduce the issue. Thanks in advance...

Subject: Re: Pentavalent carbonyl issue (urea formation?)

Posted by [padster](#) on Fri, 23 Jun 2023 18:02:09 GMT

[View Forum Message](#) <> [Reply to Message](#)

Sure thing, attached are four .dwar files with one structure each for one cycle and the .rxn file:

I'm not sure how to provide a macro to be honest

File Attachments

1) [Cycle1.dwar](#), downloaded 544 times

- 2) [Cycle2.dwar](#), downloaded 527 times
 - 3) [Cycle3.dwar](#), downloaded 516 times
 - 4) [Cycle4.dwar](#), downloaded 510 times
 - 5) [4 macrocycle disulphide.rxn](#), downloaded 537 times
-

Subject: Re: Pentavalent carbonyl issue (urea formation?)

Posted by [padster](#) on Fri, 23 Jun 2023 18:27:57 GMT

[View Forum Message](#) <> [Reply to Message](#)

Perhaps this macro works?

File Attachments

- 1) [Library_generation.dwam](#), downloaded 477 times
-

Subject: Re: Pentavalent carbonyl issue (urea formation?)

Posted by [thomas](#) on Sat, 24 Jun 2023 14:53:12 GMT





[View Forum Message](#) <> [Reply to Message](#)

Thank you for the files. To me it seems that your atom mapping is incorrect. The reactants fragments consisting out of a carbonyl group with attached bridge bond to nitrogen should not break bonding in the reaction, thus they should appear in the product the same way with the same mapping numbers. I have attached the updated macro file, which seems to work fine now.

The upper reaction shows your original mapping, the lower one shows the updated mapping:

File Attachments

- 1) [correctedMacro.dwam](#), downloaded 510 times
 - 2) [mapping.png](#), downloaded 948 times
-



Subject: Re: Pentavalent carbonyl issue (urea formation?)

Posted by [padster](#) on Sat, 24 Jun 2023 21:45:29 GMT

[View Forum Message](#) <> [Reply to Message](#)

Hi Thomas, thanks for getting back to me. However, I still get the incorrect product when I run your macro. When I try to make it myself, it rearranges the mapping so that each nitrogen is 1, 6, 9 or 12 and the first carbonyl is 2-3, but the second one is 4-5 (see attached image). It is also strange how my old mapping does not work on Windows, but it worked perfectly fine on Mac. I'm getting the idea that the Windows version is reading the mapping different or automatically changes it somehow (if that's even possible).


Image:

Structure:

File Attachments


1) [Screenshot_1.png](#), downloaded 883 times

 [Enumerate Combinatorial Library](#)



The screenshot shows a chemical reaction editor interface. At the top, there are tabs for "Generic Reaction" and "Reactants". Below the tabs is a toolbar with various icons for editing and drawing. The main workspace displays a chemical reaction scheme. The reactant is a molecule with two carbonyl groups and two nitrogen atoms. The atoms are labeled with numbers in brackets: [1-16] for the first nitrogen, [1-16] for the second nitrogen, [1-16] for the first carbonyl carbon, and [1-16] for the second carbonyl carbon. The product is a similar molecule, but the atom mapping is rearranged: the first nitrogen is labeled [1-16], the second nitrogen is labeled [12], the first carbonyl carbon is labeled [11], and the second carbonyl carbon is labeled [15]. An arrow points from the reactant to the product. Below the workspace, there is a "Use template:" dropdown menu set to "<Custom Reaction>". At the bottom left, there is a "Help" button.

- 2) [New library.dwar](#), downloaded 519 times
3) [Screenshot_2.png](#), downloaded 978 times



Subject: Re: Pentavalent carbonyl issue (urea formation?)

Posted by [thomas](#) on Sun, 25 Jun 2023 11:31:38 GMT

[View Forum Message](#) <> [Reply to Message](#)

If I import and run the macro from my last message, then I get the following product (with both versions 5.5.0 and with the current dev update):

If you use the same version on MacOS, Linux, or Windows, then it should behave exactly the same. Which version do you use anyway? The official 5.5.0, that mean the most recent (but 2 years old) official installer? Then I would recommend to update to the newest dev build (on the download page click on the 'read and understood checkbox, then click the link in the small print for you platform and download the respective archive with patch files for your installation)


Did you manually map as many atoms as it needs for auto-mapper to correctly map all remaining atoms of the reaction? If not, I suggest to remove all manual mapping and start from scratch to map at least all four O=C...N fragments manually.

Another tip: Typically, one does not use atom or bond query features on the product side of the generic reaction. Bridge bonds are an exception, because whatever matches on the reactant side needs to be reconstructed on the product side. If all your real reactants have the same bridge length, then I would remove the bridge bond from the reaction and replace it with real bonds and atoms.

One more tip: A part of the new medium sized ring comes from unknown components. For the sake of complete and perfect mapping and a stoichiometrically complete reaction you might introduce a fifth reactant with the missing atoms, that is always the same.

File Attachments

1) [correctProduct.png](#), downloaded 927 times



Subject: Re: Pentavalent carbonyl issue (urea formation?)

Posted by [padster](#) on Sun, 25 Jun 2023 18:56:21 GMT

[View Forum Message](#) <> [Reply to Message](#)

Installing the dev build seems to have done the trick! Thanks a ton!
